

# Pericyte Medium-mouse (PM-m) Catalog #1231

## **Product Description**

Pericyte Medium-mouse (PM-m), when used with Pericyte Growth Supplement (PGS-m, Cat #1282) and 10mL FBS is a complete medium designed for optimal growth of normal mouse pericytes *in vitro*. It is a sterile, liquid medium which contains essential and non-essential amino acids, vitamins, organic and inorganic compounds, hormones, growth factors, trace minerals and a low concentration of fetal bovine serum (2%). The medium is HEPES and bicarbonate buffered and has a pH of 7.4 when equilibrated in an incubator with an atmosphere of 5%  $CO_2/95\%$  air. The medium is formulated (quantitatively and qualitatively) to provide an optimally balanced nutritional environment that selectively supports the growth of normal mouse pericytes *in vitro*.

### Components

PM-m consists of 500 ml of basal medium, 10 ml of fetal bovine serum (FBS, Cat. #0010), 5 ml of Pericyte Growth Supplement-mouse (PGS-m, Cat. #1282) and 5 ml of penicillin/streptomycin solution (P/S, Cat. #0503). *Note: FBS, PGS-m and P/S are not pre-mixed in PM-m; they must be added separately to make the complete PM-m.* 

#### **Product Use**

PM-m is for research use only. It is not approved for human or animal use, or for application in *in vitro* diagnostic procedures.

### Storage

Store the basal medium at 4°C and the PGS-m, FBS and P/S solutions at -20°C. Protect from light.

### Shipping

Basal medium: room temperature. Supplements: dry ice.

#### **Instructions for Use**

Thaw PGS-m, FBS and P/S solution at 37°C. Gently tilt the tubes several times to ensure the contents are completely mixed before adding to the medium. Spray the medium bottle and tubes with 70% ethanol, and wipe to remove excess liquid. In a sterile field, remove the caps without touching the interior threads with fingers. Add PGS-m, FBS and P/S solution to the medium and mix well. Since several components are light-labile, the medium should not be exposed to light for extended periods. We do not recommend warming medium in a 37°C water bath prior to use. When stored in the dark at 4°C, the reconstituted medium is stable for one month.

*Caution: If handled improperly, some components of the medium may present a health hazard. Take appropriate precautions when handling it, including the wearing of protective clothing and eyewear. Dispose of properly.*